



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Morphology of the podocarps.—SINNOTT¹⁷ has investigated the reproductive structures of the Podocarpaceae and has come to some conclusions in reference to the relationships of the group. The detailed results are too numerous to be cited here, but the author has canvassed the structures of the ovulate strobilus, the characteristic male gametophyte, the wings of the pollen grains, the female gametophyte, fertilization, the proembryo, and the endosperm, and has concluded that the group has been derived from the Abietineae through forms resembling *Podocarpus*, which is therefore the oldest genus. Some striking resemblances between the podocarps and araucarians further suggest that both of these groups may have arisen from an ancient group closely allied to the Abietineae. A connection with the taxads is also suggested by resemblances between *Cephalotaxus* and certain species of *Podocarpus*, the conclusion being that the Taxineae, through *Cephalotaxus* (its most ancient genus), may have arisen from some ancient member of the Podocarpaceae.—J. M. C.

Adventive branches in *Frullania*.—Miss LORENZ¹⁸ finds that 4 of the 11 species of *Frullania* in New England reproduce vegetatively by adventitious shoots. From her statements it seems that a marginal cell of a leaf enlarges; that the first two planes of division are anticlinal, giving a quadrant of more or less unequal cells; that the next plane of division is periclinal; that from one of the outer cells of the resulting octant a pyramidal apical cell which gives rise to the shoot is developed. The first leaves of the shoot are rudimentary, but very soon the adult form appears. Ventral leaves are also delayed. As should perhaps be expected, these vegetative shoots are more frequent on dioecious than on autoecious species.—W. J. G. LAND.

Origin of species in *Hieracium*.—OSTENFELD¹⁹ has conducted an extensive set of experiments with *Hieracium* to discover the possible relationship between its polymorphism and its strong tendency to apogamy. He has reached the conclusion that new forms arise as hybrids and also by single variations (mutations), and that in both of these cases "the prevailing apogamy supports their existence and constancy." This means that the polymorphism of the genus is correlated with its apogamy.—J. M. C.

¹⁷ SINNOTT, EDMUND W., The morphology of the reproductive structures in the Podocarpaceae. *Ann. Botany* 27:39-82. *figs. 9. pls. 5-9.* 1913.

¹⁸ LORENZ, ANNIE, Vegetative reproduction in the New England *Frullaniae*. *Bull. Torr. Bot. Club* 39:279-284. *figs. 3.* 1912.

¹⁹ OSTENFELD, C. H., Experiments on the origin of species in the genus *Hieracium* (apogamy and hybridism). *New Phytol.* 11:347-354. 1912.